

Seasonal and Diurnal Variability of Albedo and Soil Moisture over Ranchi

Smita Dey¹ and Roshan Kumar²

¹ Ranchi University

Received: 8 December 2013 Accepted: 2 January 2014 Published: 15 January 2014

Abstract

Land surface shortwave albedo plays a central role in global and regional climate modeling. Remote sensing of surface properties and estimation of clear sky and surface albedo generally assumes that the albedo depends only on the solar zenith angle, which is verified as surface albedo decreases with increase of solar elevation angle (SAE), when SAE is greater than (value of angle=45deg.) it becomes constant. Fluctuations of soil moisture result in large variation in outgoing energy fluxes, and thus significant variation in near surface relative humidity and temperature. In this study analysis of radiation and other weather data collected from January 2009 to December 2009 at Ranchi in Jharkhand (85°30'E, 23°45'N, elev 652.272) are used to examine the diurnal and seasonal soil moisture variations with surface albedo and their relationship with soil moisture. The diurnal and seasonal soil moisture's effect from rainfall is also discussed. From the analysis of monthly data indicate that surface albedo has inverse relation with soil moisture content

Index terms— surface albedo, solar elevation angle, soil moisture.

1 Introduction

ver recent years an increase amount of attention has been paid by the atmospheric climate research community to the role of landatmospheric interactions in influencing, and being influenced by, the atmospheric structure on large scale. On these issues there are a number of national and international programs devoted. One topic which has been incorporated into these activities is the desire to come to a more comprehensive understanding of the role of surface albedo ,soil moisture ,soil thermal parameters, and there relationship in climate. It is an aspect of this general question to which this paper has been directed for Ranchi, Jharkhand, which lies at the extreme eastern end along the monsoon trough. The influence of anomalous soil moisture conditions on the atmosphere has been the subject of research for some time. Namias (1958 ??amias (, 1963)) was among the first to address the issue, noting that the seasonal anomalies of soil wetness could have impact on seasonal cycle of the atmosphere. Recently, a number of modeling studies have explicitly examined the influence of anomalies of soil moisture on the atmosphere. ??hukla and Mintz(1982) examined the impact on the atmosphere of prescribed constant anomalies of soil wetness. They demonstrated that negative anomalies of soil moisture decreases evaporation rates and increase the surface temperature. In order to regional climatic variation, soil moisture plays an very important role (Elfatih 1998 ??2008] found new algorithm, which gives a realistic estimation of soil temperature,which is helpful as the tools for interpretation of the role of heterogeneity in observed diurnal temperature variation. Soil moisture formulate the partition the available energy near land surface into sensible and latent heat exchanges with atmosphere ??Wei,1995). low pass-filter soil moisture has long memory (Pielke et al 1999;Wu et al 2002),which result in persistence of climatic anomalies. surface albedo is the fraction of incoming solar radiation reflected back to the atmosphere and space. It's a basic property of land surface and is a required component of climate and weather forecasting model [e.g, Knorr et al.,2001;Viterbo and Betts, 1999]. By influencing the absorption of solar energy, albedo helps to determine the soil heat fluxes, latent, sensible and consequently thermal and

moisture stratification of the atmospheric boundary layer. So it is imperative that the observed spatial and temporal variability in surface albedo be adequately represented in land-atmosphere models. Climate models commonly specify separate albedos for soil. Earlier research, conducted over a limited geographic area, suggested that observed spatial variability in surface albedo can be related to soil types [Tsvetsinskaya et al., 2002a; Zhou et al., 2003a]. The climate response to change in surface albedo has also been a topic of considerable study (e.g., Charney et al. 1977; Dickinson and Henderson-Sellers 1988; Xue and Shukla 1993), McCumber and Pielke (1981) performed sensitivity tests (24-h simulations) in which soil albedo was free to vary as a function of surface moisture according to Idso's formulation. The numerical experiments of Clark and Arritt [1995], who found that the albedo effect is lower than the effect of soil moisture availability for the simulation of an atmospheric convection event. Idso et al. [1975] showed in a pioneering field that bare soil is a linear function of the water content in top layer (0.2-cm to 10-cm). Such a linear relationship has been implemented in land surface model (LSMs) [Pitman et al., 1991; 2009] studied the seasonal variability in land surface albedo and soil thermal conductivity, diffusivity, soil heat capacity and their relationship with soil moisture. The result shows a typical exponential relationship as surface albedo decreases with increase of soil moisture. In dry season the thermal diffusivity increases as the power function of soil moisture. Charney (1975) discussed the effect of surface albedo's variation on the Sahara desert by the use of general circulation model. According to him change of albedo is an important factor for the formation of deserts and may be pertinent to drought conditions in Sahel. He postulates biogeophysical feedback mechanism in which lack of rainfall leading to a lack of vegetation results in a higher surface albedo. The atmosphere due to high surface albedo and positive feedback between land result in the negative effect of moisture flux convergence and rainfall, and desertification (generally in droughts). From Charney's hypothesis for the maintenance of desert, the use of more realistic albedos tends to regions of lower albedo (Cunnington and Rowntree 1986).

In the present study, A time series of continuous measurements (from Jan 2009 to Dec 2009) from the data on soil moisture, soil temperature at different depths, soil heat fluxes (at 2.56 & 5cm), air temperatures, wind speed & directions (at 1,2,4,8,16 & 32 m heights), albedo at 32 m height and all four components of radiations at 2 m height were taken from the Land Surface Atmosphere and Micrometeorological Observational System (LATAMOS) established in the Institute at Ranchi (India), which lies along extreme eastern end of monsoon trough line is used to investigate the diurnal, monthly and seasonal variations of changes in surface albedo, solar elevation angle and soil thermal parameters, and their relationship with soil moisture.

2 II.

3 Study Site And Available Data

In Ranchi annual soil temperature varies from minimum 1.0°C at surface in winter to maximum 45.0°C during summer season and soil colour varies from grey in summer to radish grey during winter with small green grass cover during monsoon period. This plateau is not only important for the evolution of monsoon trough extending from Jharkhand region along its extreme eastern end to the Rajasthan region at the western end, but also vulnerable for the thunderstorm activity during pre-monsoon summer season associated with lightning and also influence rainfall associated with lightning and cyclone during the SW monsoon period. Where the summation of incoming radiation or outgoing radiation was carried out over a specified time period. The parameters which influence surface albedo are based on the soil moisture, soil colour, solar elevation angle, roughness and so on. Solar altitude angle and soil moisture are the two main factors which influence the albedo (Li, 2009). In order to study the influence of soil moisture on the albedo, it is necessary to first examine the influence of Solar elevation angle on the surface albedo. For this, the surface albedo is calculated with Eq. (??). The Solar elevation angle can be calculated from the longitude and latitude of the site, Julian day, and mean measurement time. In addition sharp peaks in surface albedo are removed. We have taken the soil moisture data corresponding to the albedo data. For a 12 hour observation figure (2), shows that during summer Solar elevation angle is maximum and minimum for winter. From figure (3) We can see that in solar elevation influence of Solar elevation angle on surface albedo is small enough to be omitted when the solar elevation varies from 400 to 810. When solar elevation angle greater than 450, then surface albedo tends to be a constant.. In this experiment daily surface albedo calculated from solar elevation angle greater than 450 can be used to study its variation with soil moisture. The soil moisture data can also be used to calculate the surface albedo. Daily average surface albedo (?) is given by the Eq (2)

4 Global

Where is the daily average soil moisture at 15 cm. from Eq(2) decreases with increase of Remote sensing of surface properties and estimation of clear-sky and surface albedo generally assumed that the albedo variation is symmetrical about local noon, which is shown in figure ???. As the typical diurnal variation of the surface albedo on clear-day looks like U-shaped curve, in this study, the seasonal and diurnal cycle of soil moisture is observed at Ranchi, which is shown in figure ??a-d). We can see the variation of soil moisture for the four seasons. At the depth of 15 cm, 30 cm, and 45 cm Soil moisture content value is observed to be more during monsoon period when the surface is completely wet due to SW monsoon rainfall and soil sub-surface is at saturation level. The soil water content in rainy season is predominant as compare to dry season. The soil moisture at the depth of 15 cm is not maximum of all the depths. The similar variation in soil is also observed by Guan et al., (2009). the

diurnal variation reaches its minimum around 0800 LST and maximum around 1600 LST. The climatic variations of different months affect the soil moisture content. Figure (8) shows the seasonal variation of soil moisture at the depth of 5 cm, 10 cm, 20 cm, 30 cm and 50 cm for the period from January 2009 to December 2009. The moisture content is low during the dry season and it increases as the rainy season begins. The monthly average soil moisture is higher for deeper layer than the surface layer. This is may be due to the rapid evaporation of soil moisture from the surface layer.

5 III.

6 Conclusions

? The seasonal and diurnal variation of soil moisture is analyzed for the year 2009. ? It is found that the soil moisture is low during the dry season and high during the rainy season. ? The diurnal variation of surface albedo is not symmetrical about mid day. ? During sunny day, the diurnal variation of surface albedo appears as a U-shaped curve. ? The surface albedo decreases with increase of moisture content. ? When the solar elevation angle is greater than 45° then it tends to be a constant.



Figure 1: 25 Global

¹© 2014 Global Journals Inc. (US)

²© 2014 Global Journals Inc. (US) Seasonal and Diurnal Variability of Albedo and Soil Moisture over Ranchi

³© 2014 Global Journals Inc. (US)

⁴This page is intentionally left blank

⁵© 2014 Global Journals Inc. (US) -



Figure 2:




1.

Figure 3: Figure 1 :



23

Figure 4: Figure 2 :Figure 3 :




4

Figure 5: Figure 4 :




Figure 6:



5

Figure 7: Figure 5 :



297

Figure 8: - 29 GlobalFigure 7 :

Figure 9:

.1 Acknowledgements

Authors are thankful to the Dept. of Science & Technology, Govt. of India for providing support to conduct CTCZ field campaign at BIT Mesra, Ranchi.

[Water Resour. Res] , *Water Resour. Res* 34 (4) p. .

[Geophys. Res Lett] , *Geophys. Res Lett* 31 p. L11401.

[Hydrometeor] , *J Hydrometeor* . 3 p. .

[Lobell and Asner ()] , D B Lobell , G P Asner . *Moisture Effects on Soil Reflectance. Soil science society American journal* 2002. 66 p. .

[Vharney et al.] ‘1977 : A comparative study of the effect of albedo change on drought in semi-arid regions’. J G Vharney , W J Quirk , S H Chow , J Kornfield . *J. Atmos. Sci* 34 p. .

[Xue and Shukla] ‘1993: the influence of land surface properties on Sahel climate. Part 1: Desertification’. Y Xue , Shukla . *J. Climate* 6 p. .

[Elfatih and Elthahir ()] *A soil moisture-rainfall feedback mechanism 1. Theory and Observations*, A B Elfatih , Elthahir . 1998.

[Li and Hu ()] ‘A study on parameterization of surface albedo over grassland surface in the northern Tibetan Plateaus’. Y Li , Z Y Hu . 10.1007/s00376-009-0161-6. *Adv. Atmos. Sci* 2009. 26 (1) p. .

[Lawrence and Slingo ()] ‘An annual cycle of vegetation in a GCM. Part 1: implementation and impact on evaporation’. D M Lawrence , J M Slingo . 10.1007/s00382-003-0366-9. *Clim. Dyn* 2004. 22 p. .

[Tsvetsinskaya et al. ()] ‘An integrated assessment of landscape evolution, long-term climate variability, and use in the Amudarya Prisyrykamysh delta’. E A Tsvetsinskaya , B I Vainberg , E V Glushko . *J. Arid Environ* 2002b. 51 (3) p. .

[Nai et al. ()] *common land model (CLM) TEchnicals Documentation and User’s Guide*, Y X Nai , R E Zeng , Dickinson . 2001.

[Zhou ()] ‘comparison of seasonal and spatial variations of albedos from Moderate-Resolution Imaging Spectroradiometer (MODIS) and Common Land model’. L Zhou . doi:10.1029. *J. Geophys. Res* 2003a. 2002 JD003326. 108 (D15) p. 4488.

[Gao et al. ()] ‘Comparison of two soil temperature algorithms for a bare ground site on the Loess Plateau in China’. Z Q Gao , D H Lenschow , R Horton , M Y Zhon , L L Wang , J Wen . *J. Geophys. Res* 2008. 113 p. D18105.

[Matsui et al. ()] ‘Continentscale multiobservation calibration and assessment of Colorado state University Unified Land Model by application of Moderate resolution Imaging spectroradiometer (MODIS) surface albedo’. T Matsui , A Beltrán-Przekurat , R A Pielke , D Niyogi , M B Coughenour . doi: 10. 1029. *J. Geophys.Res* 2009. 2006J G000229. 112 p. 2028.

[Duke and Guérif ()] ‘Crop reflectance estimate errors from the sail model due to spatial and temporal variability of canopy and soil characteristics’. C Duke , M Guérif . doi:10.1016 / S0034-4257(98)0. *Remote Sens. Environ* 1998. 66 (3) p. .

[Pitman et al. ()] ‘Description of the bare essentials of surface transefer for the bureau of’. A J Pitman , Z L Yang , J G Cogley , A Henderson-Sellers . *Meteorol.Res.Centre AGCM,BMRC Res. Report* 1991. (32) .

[Timbal et al. ()] ‘Does soil moisture influence climate variability and predictability over Australia?’. B Timbal , S Power , R Columan , J Vivian , S Lirola . *J.Climate* 2002. 15 p. .

[Charney ()] ‘Dynamics of deserts and drought in the Sahel’. J Charney . *Quart. J. Roy . Meteor. Soc* 1975. 101 (428) p. .

[Hoffer and Johannsen ()] *Ecological potential in spectral signatures analysis. Remote Sensing in Ecology*, R M Hoffer , C J Johannsen . 1969. Athens (GA. p. . University of Georgia

[Viterbo and Betts ()] ‘Impact on ECMWF forecasts of change to the albedo of the boreal forest in presence of snow’. P Viterbo , A K Betts . *J. Geophys. Res* 1999. 104 (D22) p. .

[Shi ()] *Initial soil moisture effects on the climate in China (Oceanic and Coastal Sea Research*, X Shi . 2009. 8 p. .

[Dickinson and Henderson-Sellers ()] ‘Modeling tropical deforestation: A study of GCM land-surface parameterizations’. R E Dickinson , A Henderson-Sellers . *Quart. J. Roy. Meteor. Soc* 1988. 114 p. .

[Clark and Arritt ()] ‘Numerical Simulations of the Effect of Soil Moisture and Vegetation cover on the Development of Deep Convection’. C A Clark , P W Arritt . 10.1175/15200450. *J. Appl. Meteorol* 1995. 1995. 34 p. . (034 ;2029 : NSO TEO;2.0.CO;2.)

[Namias (ed.) ()] *Persistence of mid-tropospheric circulation between adjacent months and seasons. The Atmospheric and Sea in Motion*, J Namias . P.Polin (ed.) 1958. Rockefeller Institute Press and Oxford University Press. p. .

- [Koster ()] ‘Regions of strong coupling between soil moisture and precipitation’. R Koster . *Science* 2004. 305 p.
- [Liu et al. ()] ‘Relationships between surface albedo, soil thermal parameters and soil moisture in the semi-arid area of Tongyu, north-easterns China’. H Z Liu , B M Wang , C B Fu . 10.1007/s00376-008-0757-2. *Adv. Atmos. Sci* 2008. 25 (5) p. .
- [Douville and Chauvin ()] ‘Relevance of soil moisture for seasonal climate predication: A preliminary study’. H Douville , F Chauvin . *Clim. Dyn* 2000. 16 p. .
- [Liu et al. ()] ‘Rwlatng soil surface moisture to reflectance’. W D Liu , F Baret , X F Gu , Q X Tong , L F Zheng , B Zhang . 10.1016/S0034-4257(01)00347-9. *Remote Sens. Environ* 2002. 81 p. .
- [Pielke et al. ()] ‘Seasonal weather predication as an initial value problem’. R A Pielke , G E Liston , J Eastman , X Lu . *J. Geophys. Res* 1999. 104 p. 479.
- [Gascoin et al. ()] ‘Sensitivity of bare soil albedo to soil moisture on the Zongo glacier (Bolivia)’. S Gascoin , A Ducharne , P Ribstein , E Perroy , P Wang . doi: 10. 1029. *Geophys. Res. Lett* 2009a. 2008. 36 p. L036377.
- [Mc Cumber and Pielke ()] ‘Simulation of the effect of surface fluxes of heat and moisture in a mesoscale numerical model 1. Soil layer’. M C Mc Cumber , R A Pielke . *J. Geophys. Res* 1981. 86 p. .
- [Cunnington and Rowntree ()] ‘Simulations of Saharan atmosphere -dependence of moisture and albedo’. W M Cunnington , P R Rowntree . *Quart.J. Roy. Meteor. Soc* 1986. 112 p. .
- [Wei ()] ‘Soil moisture’. M Y Wei . *NASA Conference Publication 3319*, (Tiburón, California) 1995. p. 80. (Report)
- [Lakshmi et al. ()] *Soil moisture as an indicator of weather extremes*, V Lakshmi , T Piechota , U Narayan , C Tang . 2004.
- [Idso et al. ()] ‘The dependence of bare soils albedo on soil water content’. S Idso , R Jackson , R Reginato , B Kimball , F Nakayama . 10.1175/1520-0450(1975)014;0109:TDOBSA;2.0.CO;2. *J. Appl. Meteorol* 1975. 14 p. .
- [Shukla and Mintz ()] ‘The influence of land surface evoparation on Earth’s climate’. J Shukla , Y Mintz . *Science* 1982. 215 p. .
- [Wu et al. ()] *The response of soil moisture to long-term variability of precipitation*, W R Wu , M A Geller , R E Dickinson . 2002.
- [Knorr et al. ()] ‘The role of bright desert regions in shaping North African climate’. W Knorr , K.-G Schnitzler , Y Govaerts . *Geophys. Res. Lett* 2001. 28 (18) p. .
- [Guan et al. ()] ‘Variability of soil moisture and its relationship with surface albedo and soil thermal parameters over the Loess Plateau’. X D Guan , J P Huang , N Guo , J Bi , G Wang . *Adv. Atmos. Sci* 2009. 26 (4) p. .
- [Wang et al. ()] ‘Variation of surface albedo and soil thermal parameters with soil moisture content at a semi-desert site on the western Tibetan plateau’. K C Wang , P C Wang , J M Liu , M Sparrow , S Haginoya , X J Zhon . 10.1007/s10546-004-7403-z. *Bound-Lay. Meteorol* 2005. 116 p. .